

The effects of facility dogs on burnout, job-related well-being, and mental health in pediatric hospital professionals

Clare L. Jensen¹, Jessica Bibbo^{1,2}, Kerri E. Rodriguez^{1,3}, and Marguerite E. O’Haire¹

¹Center for the Human-Animal Bond, Department of Comparative Pathobiology, College of Veterinary Medicine, Purdue University, West Lafayette, IN, USA

²Center for Research and Education, Benjamin Rose Institute on Aging, Cleveland, OH, USA

³Human-Animal Bond in Colorado, School of Social Work, Colorado State University, Fort Collins, CO, USA

Correspondence concerning this article should be addressed to Clare L. Jensen, Center for the Human-Animal Bond, Purdue University College of Veterinary Medicine, 725 Harrison Street, West Lafayette, IN 47907. Email: jensen76@purdue.edu. Telephone: 317-603-1379.

Additional Author Email Addresses: jbibbo@benrose.org (Jessica Bibbo), kerri.rodriguez@colostate.edu (Kerri Rodriguez), mohaire@purdue.edu (Marguerite O’Haire)

Conflicts of Interest: None.

Funding: This research was supported by a grant from Elanco Animal Health, a division of Eli Lilly and Company. This publication was made possible with partial support from Grant #KL2TR001106 and Grant #UL1TR002529 (A. Shekhar, PI), from the National Institutes of Health, National Center for Advancing Translational Sciences, Clinical and Translational Sciences Award.

Acknowledgements: The authors sincerely appreciate Katelynn Burgess for her lead role in data collection and all participants for the time spent sharing their experiences. We would also like to acknowledge Canine Assistance with gratitude for their participation in this study.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/JOCN.15694](https://doi.org/10.1111/JOCN.15694)

This article is protected by copyright. All rights reserved

MS. CLARE JENSEN (Orcid ID : 0000-0002-4664-5774)

Article type : Original Article

Abstract

Aims and objectives

The present study aimed to examine the effect of working with a facility dog on pediatric healthcare professionals' work-related burnout, job perceptions, and mental health.

Background

Due to their roles caring for ill children and distressed families, pediatric healthcare professionals often experience substantial depression and burnout. According to prior research, facility dogs in children's hospitals may provide significant benefits to pediatric patients. However, their potential effects on healthcare professionals have been minimally explored.

Design

A cross-sectional design was used in adherence to the STROBE checklist.

Method

Among 130 participants, $n=65$ pediatric healthcare professionals working with a facility dog were compared to $n=65$ control participants matched on age, gender identity, job position category, and pet ownership. Hierarchical regression assessed the effect of working with a facility dog on standardized self-report measures of work-related burnout, job perceptions, and mental health.

Results

For work-related burnout, working with a facility dog was associated with higher perceived personal accomplishment, but had no effect on emotional exhaustion. With respect to job perceptions, working with a facility dog was associated with more positive job descriptions and lower intention to quit, but not with perceptions about co-workers or workplace social support. Finally, in relation to mental health, working with a facility dog was associated with more

positive emotions, better perceived mental health, and less depression, but had no effect on anxiety.

Conclusion

Findings suggest that facility dogs may be related to several benefits for healthcare professionals' work-related burnout, job perceptions, and mental health, but that they do not influence all components of these areas.

Relevance to clinical practice

The present research functions to inform personnel in pediatric hospitals with existing facility dog programs on the scope of their effects, in addition to shaping the expectations of hospitals considering the addition of a facility dog program.

Keywords: Burnout, Professional; Hospitals, Pediatric; Personnel Turnover; Mental Health; Job Satisfaction; Hospital, Nursing Staff; Animal Assisted Therapy; Dogs

'What does this paper contribute to the wider global clinical community?'

- According to prior research, facility dogs, or specially trained full-time therapy dogs, can benefit hospitalized children and their families. The present research suggests that facility dogs are also benefitting the healthcare professionals they work alongside.
- Compared to a matched control group, pediatric healthcare professionals working with a facility dog reported less work-related burnout, better job-perceptions, and better functioning in most areas of mental health.
- Results provide preliminary evidence of an intervention that may enhance healthcare professionals' well-being while also setting realistic expectations for both existing and future facility dog programs.

Introduction

There is a growing practice of incorporating therapy dogs into daily patient care, especially in pediatric hospitals. According to prior research, therapy dogs may provide significant benefits to pediatric patients' physical and mental health, including reduced fear, anxiety, and perceived pain (e.g., Crossman et al., 2020; Lindström Nilsson et al., 2020). Some hospitals feature full-time therapy dogs, termed "facility dogs," that work with pediatric healthcare professionals in goal-directed practices to improve the patient experience (Shilp, 2019).

Facility dogs have become increasingly common to fill unique roles alongside professionals in careers of service for their communities and the public (e.g., social workers, school counselors, forensic interviewers, healthcare professionals; Shilp, 2019). These dogs are distinguished from therapy dogs in that they undergo extensive and rigorous training similar to that of a service dog. However, unlike a service dog's training for disability-related tasks to assist a specific individual with whom they are paired, facility dogs are trained to assist a number of community members with a range of needs (i.e., the clients or patients of the professional they are working alongside; Walsh et al., 2018). Workplaces that incorporate facility dogs include hospitals, medical centers, educational settings, police stations, and courthouses (e.g., Spruin et al., 2020; Barnett, 2019; Shilp, 2019). Depending on the profession in which they will be engaged, facility dogs might be taught skills such as remaining discrete in court, resting their chin on a person's lap, or moving delicately around medical equipment. In children's hospitals with facility dog programs, these dogs are regularly present to augment medical interventions and interactions by offering distraction, comfort, and motivation to patients.

While the primary purpose of many hospital facility dog programs is to benefit the patients, an equally important population in contact with facility dogs is the team of healthcare professionals they work alongside. Healthcare professionals play a crucial role in the human experience and have great potential to influence the lives of many. However, healthcare professionals often experience chronic stress that puts them at risk for work-related burnout and poor mental health (Al Sabei et al., 2020; Ortega-Campos et al., 2019; Tawfik et al., 2018). Work-related burnout may lead to adverse effects spanning multiple facets of intra- and inter-personal health and well-being. On a community level, burnout can have detrimental effects on professional and hospital functioning, such as inadequate job-performance (Lee et al., 2019), increased errors with detriments to patients, and additional costs on the hospital due to

absenteeism and turnover (Buckley et al., 2020). The development of strategies for reducing burnout is vital to the optimization of hospital financial viability and patient care quality (Wei et al., 2020). For pediatric healthcare professionals specifically, evidence is limited regarding interventions to effectively reduce burnout and improve well-being (Buckley et al., 2020).

Prior research aimed at reducing these negative mental health outcomes among healthcare professionals has found that social support may play a critical role in buffering the adverse effects of burnout (Velando- Soriano et al., 2020). However, meaningful social support is not limited to human sources as emerging human-animal interaction research suggests that social support from dogs may have similar benefits. As it is applied to human-animal interactions, the social support theory suggests that dogs may be a unique source of non-judgmental companionship (Beetz, 2017). Additionally, the presence of a dog may result in more frequent and positive social exchanges with other people in one's environment (Wood et al., 2007). In fact, dogs have been found to assist in facilitating conversations and establishing inter-personal connections (Beetz, 2017; Wells, 2019). Studies have found that adults with pet dogs report greater well-being compared to people without pet dogs, including less loneliness (Carr et al., 2020; Powell et al., 2019) and lower risk for depression after a social loss (Carr et al., 2020). There is also evidence that the social support provided by dogs may function as a mechanism for buffering stress (Beetz, 2017; Ein et al., 2018). In one pioneering study, female participants undergoing a stressful task showed less stress reactivity with their pet dog present than with a close friend present (Allen et al., 1991). This set the stage for continued research on the stress-reducing effects of social support from dogs, which has found evidence that the presence of an unfamiliar therapy dog may also confer stress buffering benefits (e.g., S. B. Barker et al., 2010; Polheber & Matchock, 2014).

In addition to their general stress-reducing effects, dogs may also affect stress and well-being in the workplace. In organizations that allow pet dogs, research has found that dog presence is associated with lower physiological and perceived stress at work (R. T. Barker et al., 2012; Hall et al., 2017). Dogs in the workplace have also been related to more positive perceptions of the work environment, increased communication among coworkers (R. T. Barker et al., 2012; Hall et al., 2017), and improved morale (Foreman et al., 2017). These observations of the benefits of dogs in the workplace have also been investigated specifically within the healthcare setting. A survey conducted before and after the implementation of a therapy dog

visitation program indicated that healthcare professionals reported a more positive work environment and improved clinic climate after program implementation (Yordy et al., 2020). Further, a recent randomized trial found that emergency medicine healthcare professionals experienced lower self-reported stress after a five-minute break with a therapy dog compared to coloring (Kline et al., 2020). In an earlier study, it was demonstrated that just five minutes with a therapy dog could produce the same level of stress-reduction among healthcare professionals as 20 minutes of quiet rest (S. B. Barker et al., 2005).

Despite promising initial findings on therapy dog interactions for reducing momentary stress, to our knowledge, there are no published empirical studies on the broader effects of full-time facility dogs working with pediatric healthcare professionals. Thus, to begin filling this gap, the present study seeks to quantify the effects of working with a facility dog on several key areas of pediatric hospital professional functioning. We hypothesized that pediatric healthcare professionals working with facility dogs would report less work-related burnout, more positive job perceptions, and better mental health than matched control participants working without facility dogs.

Method

Participants

The study sample included $N=130$ pediatric healthcare professionals working in children's hospitals across the United States. Participants either worked directly with a facility dog (treatment group; $n=65$), or did not work with a facility dog (control group; $n=65$). Inclusion criteria for all participants included over 18 years of age and currently employed in a pediatric hospital. For the facility dog group, inclusion criteria consisted of working with a facility dog for at least six months prior to study participation. Participants were paired across groups based on four key matching characteristics. These characteristics included age (± 10 years), gender identity, pet ownership, and job title category. Initial matching efforts also included number of years working in the profession (± 5 years), but this was later dropped as a matching criteria to recruit the final set of control participants ($n=122$, 94% of participants were matched on number of years working in the job position).

Sample demographics are reported in Table 1. The vast majority of participants identified as female (92%) and white (95%). The mean age was 37 years ($SD = 10.4$) and the mean time in

present job position was 5 years and 8 months ($SD = 5$ years and 7 months). Ten position categories were used to describe participants' job titles, with most participants working in child-life (35%) or administration/managerial positions (38%), in addition to nurses, social workers, chaplains, physicians, therapists, psychologists, nursing-administration, and child-life-administration. The majority of participants were married or living with a significant other (64%), had children (55%), and had one or more pet dogs at home (65%).

All facility dogs were raised, trained, and placed by the organization Canine Assistants, a nonprofit service and facility dog provider located in Alpharetta, Georgia. Dogs were bred on-site at Canine Assistants and included Golden Retrievers, Labrador Retrievers, Goldendoodles, and mixes of the three. After an average of 18 months spent learning their specific skill-sets, facility dogs were eligible to be partnered with pediatric healthcare professionals. At least six months prior to study participation, pediatric healthcare professionals who had applied and been approved for the Canine Assistants program traveled to the organization's campus for a two-week partnering session. During these partnering sessions, healthcare professionals were matched with a facility dog, guided through exercises in bond-building, and taught how best to work with their facility dogs in healthcare settings.

Procedure

The research protocol was approved by the Purdue University Human Research Protection Program Institutional Review Board (IRB Protocol 1607017967) and by the Purdue Animal Care and Use Committee (PACUC Protocol 1702001541). The present study adhered to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement and checklist (von Elm et al., 2008; Supplementary File 1).

Potential participants were recruited with an initial mailed study packet that contained information about the study and \$10 as remuneration for reviewing the study details. A member of the research team then called potential participants to answer any questions about the study and obtain voluntary verbal informed consent. Consented participants then received an email with a link to complete an online survey, in which they verified their informed consent electronically prior to the first survey item. Participants were informed that their individual answers were confidential and that their responses would not be shared with Canine Assistants, their employers, or anyone outside of the research team.

Participant recruitment was conducted in three phases. During phase one, individuals were recruited directly from the provider Canine Assistants, including those who were previously placed with a facility dog or on the waitlist to receive a facility dog. In phase two, participants with a facility dog were asked to recommend people working *without* a facility dog who were otherwise similar to themselves on the research team's key matching characteristics. During phase three of recruitment, a list was compiled with the characteristics required to complete matching for any remaining participants in the treatment group. Information was provided in ranges determined to sufficiently prevent the identification of any participants. This list and the study flyer were distributed widely to prior participants, affiliates of the research team and facility dog provider, and attendees of a national conference for healthcare professionals with facility dogs.

As part of a larger two-week study period (involving daily ecological momentary assessment), participants were asked to fill out an online survey with the opportunity for compensation of \$45. In total, 219 people (82%) consented to participate in the study. Of those who consented, 147 (67%) were eligible to participate. Of those who were eligible, 139 (95%) completed the study and 130 (88%) could be matched with another participant across groups. Data was collected between February 2018 and March 2019.

Measures

Participants completed the online survey via the website Qualtrics. Survey items included demographic characteristics (Table 1), pet and facility dog (when applicable) descriptor items, open-ended questions, and standardized measures of work-related burnout, job perceptions, and mental health. Number of items, scoring, and Cronbach's α (i.e., reliability and internal consistency within the measure) are listed for each measure in Table 2.

Work-Related Burnout

Maslach Burnout Inventory (MBI): Emotional Exhaustion

On the MBI subscale for emotional exhaustion (Maslach et al., 1997), each item asks participants to indicate how often they experience the given statement on a scale of zero (never) to six (every day). Items are reverse scored such that higher scores indicate a better outcome (i.e., less emotional exhaustion). Both sub-scales of the MBI have strong psychometric properties and have been widely used in healthcare professions.

Maslach Burnout Inventory (MBI): Personal Accomplishment

The MBI subscale for personal accomplishment (Maslach et al., 1997) also asks participants to indicate how often they experience the given statement for each item, using the same scale of zero (never) to six (every day). Higher scores indicate greater perceptions of personal accomplishment.

Job Perceptions**Job Related Depression-Enthusiasm Scale (JRDES)**

The JRDES (Warr, 1990) is a theoretically-based and validated measure (Laguna et al., 2017), designed to assess mental well-being directly related to the participant's job. Participants are asked to rate from one (never) to five (all of the time), for the past few weeks, how much of the time their job has made them feel each of the listed emotions. Higher scores indicate better outcomes (i.e. lower job-related depression and higher job-related enthusiasm).

Job Descriptive Index (JDI): People on Your Present Job

The JDI was used to assess perceived interpersonal quality of the work environment via the People on Your Present Job subscale (Smith et al., 1969). Each item asks participants to indicate whether a given adjective describes the people they work with. Response options are "Yes," "No," and "?", with "?" always being scored as one. In positively worded items, "Yes" is scored as three and "No" is scored as zero, whereas in negatively worded items, "Yes" is scored as zero and "No" is scored as three. Higher scores indicate more positive descriptions of people at the job.

Job in General (JIG)

Designed to accompany subscales of the JDI, the Job in General (JIG) scale is a global measure of how people feel about their jobs the majority of the time (Ironson et al., 1989). Each item asks participants to indicate whether a given adjective describes their job. The JIG uses the same response options and is scored in the same manner as the JDI subscale above. Higher scores indicate more positive global descriptions of one's job.

Workplace Social Support

Accepted Article

Perceived social support at work was assessed as an average of coworker support, supervisor support, and organizational support (3 items each; Yoon & Thye, 2000). Participants are asked to select how much they agree or disagree with statements ranging from one (strongly disagree) to five (strongly agree). Higher average scores indicate greater perceived support.

Anticipated Turnover Scale (ATS)

The ATS quantifies a person's intention to voluntarily terminate their current employment (Mowday et al., 1984). This scale has been used extensively with healthcare professionals (Kim & Kao, 2014) and it has good validity and reliability. Each item provides the participant with a statement about the possibility of quitting their current position and asks them to indicate how much they agree with the statement. Higher scores indicate less possibility of quitting one's job position.

Turnover Intention Scale (TIS-6)

The TIS-6 is a concise measure of participants' plans to stay in their current job position (Bothma & Roodt, 2013). Each item asks the participant to indicate to which degree they experience the listed thought or feeling about staying or leaving their job. Higher scores indicating greater intention to leave.

Mental Health

PROMIS Short Form v1.0: Depression and Anxiety

The Patient-Reported Outcomes Measurement Information System (PROMIS) was developed by the National Institute of Health (NIH) as a precise and accurate measurement of patient-reported health status for physical, mental, and social well-being to be compared across studies (Cella et al., 2010). PROMIS measures in the present study included Depression v1.0 adult short form (8a) and Anxiety v1.0 adult short form (8a) administered in English via unassisted online self-report. Items in both instruments ask the participant to rate how often they experienced the listed feeling in the prior seven days, with response options on a scale of one (never) to five (always). Raw scores were converted to measure-specific T-scores using the conversion tables published in the PROMIS Adult Profile Scoring Manual with a population mean of 50 and standard

Accepted Article

deviation of 10. Higher scores indicate more anxiety or worse depression, respectively. Both short-form instruments were found to have superior reliability and validity (Schalet et al., 2016).

PROMIS Global Health v1.2: Mental Health

Another PROMIS measure, the PROMIS Scale v1.2-Global Mental Health adult short form (2a), was included to assess participants' overall perceptions of their mental health. This scale first asks participants to rate their general mental health, mood, and ability to think, followed by their general satisfaction with social activities and relationships. Response option categories range from one (excellent) to five (poor). Raw scores were converted to measure-specific T-scores using the conversion tables published in the PROMIS Global Health Scoring Manual. Higher scores represent worse mental and social health. This measure of mental health has been found to have good validity and reliability (Hays et al., 2017).

Scale of Positive and Negative Experience (SPANE)

This validated scale was used to assess a participant's socioemotional well-being and affect range, with sub-scales for positive and negative feelings (Diener et al., 2010). Items ask the participant to rate how often they experienced the listed feeling over the past 4-week period, on a scale of one (very rarely or never) to five (very often or always). Higher scores indicate greater valence of emotion (i.e., more positive or more negative). Affect balance score was calculated by subtracting negative affect score from the positive affect score.

Data Analysis

A power calculation was not conducted a priori due to the convenience sampling required of this study, wherein there was a fixed maximum sample size in the study population (i.e., working with one facility dog provider). Thus, we recruited as many participants as possible until the population available had been exhausted. To determine post-hoc power, we conducted a power analysis to detect a medium effect size ($d=0.5$), which yielded a sample size of $N=120$ across two groups ($n=60$ per group). Preliminary analyses were conducted to test for demographic differences between groups using t-tests for continuous variables and chi-squared

Accepted Article

for categorical variables. Due to the racial and ethnic homogeneity of the sample, comparisons could not be made between specific racial and ethnic categories. However, based on the role that minority status may play in mental health and job-related well-being (Boateng et al., 2019), this variable was not excluded and was recoded into a dichotomous variable for minority (Hispanic or Latino, Black or African American, Asian Indian, American Indian or Alaska Native; when selected in any combination with or without the category White) or non-minority (White; when selected as the only racial/ethnic category) status. Relationship status was similarly recoded into a dichotomous variable for single (single, never married/divorced/separated/widowed) or in a relationship (married/living with my partner or significant other, not married).

A series of hierarchical regression models were conducted to determine the effect of working with a facility dog on work-related burnout, job perceptions, and mental health. Models were conducted with all potential explanatory covariates included to account for as much explained variance in mental health and job-related outcomes as possible. Continuous demographic covariates included age, number of children, and time in job position. Dichotomous (1/0) demographic covariates included gender identity, relationship status, pet dog ownership, and minority racial/ethnic status. All demographic variables were entered in the first step of models, while study group (facility dog or control group) was entered as a second step. Dependent variables were survey measure scores calculated at the scale and sub-scale levels.

From each model, R^2 was used to estimate the proportion of outcome variance explained by the included independent variables. Thus, the R^2 change between the first and second model for each outcome was used to estimate the proportion of variance uniquely explained by the addition of the facility dog, over and above what could be explained by demographic variables. Subsequent F-statistics and p-values, from the corresponding F-tests for each model, were used to determine whether R^2 values for proportion of explained variance were significant. If the F-change was significant, the addition of the facility dog in the second model significantly improved the proportion of variance explained for that outcome. Taken together, where the outcome (i.e., measures of work-related burnout, job perceptions, mental health) was different between the control group and facility dog group, F-statistics were used to indicate what that difference was related to. The first model indicated how much of the outcome difference was related to participant demographics. The second model indicated, beyond what was related to demographics, how much of the outcome difference was related to the facility dog.

Effect sizes were calculated as Cohen's d values, with the cutoff values of 0.2 for small, 0.5 for medium, and 0.8 for large effects (Cohen, 1992). Analyses were conducted using IBM SPSS Statistics Version 26.

Results

Preliminary Analyses

Group-specific descriptive statistics for demographic covariates are shown in Table 1. Due to participant matching, there were no significant differences in age, gender identity, pet ownership status, and job position category across groups. Although a small portion of the sample ($n=8$) had not been matched for time in job position, the majority of the sample (94%) was matched on this characteristic and a t-test showed no significant difference between groups. There were no significant differences between groups in the unmatched characteristics of race/ethnicity, relationship status, children, or dog-specific pet ownership. Correlations between covariates were found to be primarily weak ($r \leq .30$), with a moderate correlation between age and time in job position ($r = .60$). Analyses were conducted to test the assumptions of normality, linearity, multicollinearity, and homoscedasticity for each model. All assumptions were satisfied, except for the assumption of normality for the JDI and JIG measures. Because the present sample size provided adequate robustness to the test (Schmidt & Finan, 2018), authors proceeded without meeting the assumption of normality for JDI and JIG measures. Implications of this are further discussed in the limitations. Group mean comparisons and Cohen's d effect sizes for each measure are shown in Table 2. Hierarchical regression model results are shown in Table 3.

Work-Related Burnout

Maslach Burnout Inventory: Emotional Exhaustion

In regression model one, demographic characteristics explained a significant amount of variance in emotional exhaustion ($p < .001$). The addition of facility dog presence in model two made no significant contribution ($\Delta R^2 = 0.00$). There was no significant association between working with a facility dog and emotional exhaustion. Facility dog presence or absence was only related to a 0.05 standard deviation difference in model adjusted means, and the effect size of this difference was minimal ($\beta = 0.05, p = .513, d = 0.08$).

Maslach Burnout Inventory: Personal Accomplishment

The variance in personal accomplishment was not significantly explained by demographic characteristics in regression model one ($p = .955$). After the addition of facility dog presence in model two, there was a significant increase in explained variance with a 16% change in the coefficient of determination ($\Delta R^2 = 0.16$). Results showed a significant association between facility dog presence and personal accomplishment, such that participants working with a facility dog reported greater perceived accomplishment at work than participants in the control group. Compared to the control group, participants with a facility dog scored 0.42 standard deviations higher for personal accomplishment, with a large effect size ($\beta = 0.42, p < .001, d = 0.91$).

Job Perceptions

Job Related Depression-Enthusiasm Scale (JRDES)

Demographic characteristics in model one significantly explained a portion of the variance in job-related depression and enthusiasm ($p = .016$). Explained variance was increased by the addition of facility dog presence in model two with a 6% change in the coefficient of determination ($\Delta R^2 = 0.06$). Thus, working with a facility dog was a significant predictor of greater job-related enthusiasm and less job-related depression, such that participants with a facility dog scored 0.24 standard deviations higher than the control group with a moderate effect size ($\beta = 0.24, p = .005, d = 0.48$).

Job Descriptive Index (JDI): People on Your Present Job

Variance in the JDI subscale for perceptions of people at participants' jobs was not explained by demographic characteristics in model one ($p = .188$) and the addition of facility dog presence in model two did not show a significant change ($\Delta R^2 = 0.01$). There was no significant association between working with a facility dog and perceptions about people at work. Comparing facility dog and control groups, the model adjusted mean difference was only 0.12 standard deviations and the effect size was small ($\beta = 0.12, p = .197, d = 0.22$).

Job in General (JIG)

Demographic characteristics in model one significantly explained a portion of the variance in the JIG scale for participants' perceptions of their jobs overall ($p = .032$). Explained variance was increased by the addition of facility dog presence in model two with a 6% change in the coefficient of determination ($\Delta R^2 = 0.06$). Thus, working with a facility dog was a significant predictor of better perceptions about the job overall, with a group difference of 0.25 standard deviations and a large effect size ($\beta = 0.25, p = .004, d = 0.57$).

Workplace Social Support

Variance in participants' perceived social support at work was not explained by demographic characteristics in model one ($p = .057$) and the addition of facility dog presence in model two did not show a significant change ($\Delta R^2 = 0.02$). There was no significant association between working with a facility dog and perceived social support at work. Although effect size was moderate, the model adjusted mean difference was only 0.16 standard deviations ($\beta = 0.16, p = .074, d = 0.31$).

Anticipated Turnover Scale (ATS)

The variance in participants' plans to leave their job was not significantly explained by demographic characteristics in regression model one ($p = .232$). After the addition of facility dog presence in model two, there was a significant increase in explained variance with a 6% change in the coefficient of determination ($\Delta R^2 = 0.06$). Results showed a significant association between facility dog presence and anticipated turnover, such that participants working with a facility dog reported less anticipation of leaving their jobs than participants in the control group. With a group difference of 0.24 standard deviations, there was a moderate effect size ($\beta = 0.24, p = .006, d = 0.47$).

Turnover Intention Scale (TIS-6)

The variance in participants' intentions to quit their jobs was not significantly explained by demographic characteristics in regression model one ($p = .131$). After the addition of facility dog presence in model two, there was a significant increase in explained variance with a 7% change in the coefficient of determination ($\Delta R^2 = 0.07$). Results showed a significant association between facility dog presence and turnover intention, such that

Accepted Article

participants working with a facility dog reported less intentions to quit their jobs than participants in the control group. Compared to the control group, participants with a facility dog scored 0.27 standard deviations lower for turnover intention, with a large effect size ($\beta = -0.27, p = .002, d = -0.50$).

Mental Health

PROMIS: Depression

Demographic characteristics in model one significantly explained a portion of the variance in depression ($p = .051$). Explained variance was increased by the addition of facility dog presence in model two with a 4% change in the coefficient of determination ($\Delta R^2 = 0.04$). Thus, working with a facility dog was a significant predictor of less depression, such that participants with a facility dog scored 0.20 standard deviations lower than the control group with a moderate effect size ($\beta = -0.20, p = .025, d = -0.40$).

PROMIS: Anxiety

In regression model one, demographic characteristics explained a significant amount of variance in anxiety ($p = .014$). The addition of facility dog presence in model two made no significant contribution ($\Delta R^2 = 0.01$). There was no significant association between working with a facility dog and less anxiety. Comparing facility dog and control groups, the model adjusted mean difference was only 0.12 standard deviations and the effect size was small ($\beta = -0.12, p = .186, d = -0.20$).

PROMIS: Mental Health

Demographic characteristics in model one significantly explained a portion of the variance in overall mental health ($p = .011$). Explained variance was increased by the addition of facility dog presence in model two with a 4% change in the coefficient of determination ($\Delta R^2 = 0.04$). Working with a facility dog was a significant predictor of better overall mental health, with a group difference of 0.21 standard deviations and a moderate effect size ($\beta = -0.21, p = .017, d = -0.47$).

Scale of Positive and Negative Experience (SPANE): Positive Affect

Demographic characteristics in model one significantly explained a portion of the variance in positive affect ($p = .001$). Explained variance was increased by the addition of

Accepted Article

facility dog presence in model two with a 8% change in the coefficient of determination ($\Delta R^2 = 0.08$). Thus, working with a facility dog was a significant predictor of more positive affect. Facility dog presence was associated with greater positive affect, with a difference of 0.29 standard deviations and a large effect size ($\beta = 0.29, p < .001, d = 0.62$).

Scale of Positive and Negative Experience (SPANE): Negative Affect

Demographic characteristics in model one significantly explained a portion of the variance in negative affect as well ($p = .001$). Explained variance was increased by the addition of facility dog presence in model two with a 3% change in the coefficient of determination ($\Delta R^2 = 0.03$). Thus, working with a facility dog was also a significant predictor of less negative affect, with a group difference of 0.18 standard deviations and a moderate effect size ($\beta = -0.18, p = .031, d = -0.30$).

Scale of Positive and Negative Experience (SPANE): Balanced Affect

On the final measure, demographic characteristics in model one significantly explained a portion of the variance in affect balance ($p < .001$). Explained variance was increased by the addition of facility dog presence in model two with a 7% change in the coefficient of determination ($\Delta R^2 = 0.07$). Working with a facility dog was a significant predictor of better affect balance, such that groups differed by 0.27 standard deviations with a large effect size ($\beta = 0.27, p = .001, d = 0.53$).

Discussion

The goal of the present study was to evaluate the effects of facility dogs on pediatric healthcare professionals' work-related burnout, job perceptions, and mental health. A cross-sectional design compared a treatment group who worked with facility dogs to a matched control group using standardized self-report measures. Findings were mixed, with specific areas showing significant relationships with working with a facility dog and others showing no differences between groups. The nuanced variances within these study findings indicate facility dogs may have differing effects on specific aspects of job-related and overall well-being.

Within the framework of work-related burnout, working with a facility dog was associated with greater feelings of personal accomplishment at work. This is consistent with quotes from

Accepted Article

facility dog handlers describing how rewarding it can be to see patients and families benefitting from the facility dog (Barnett, 2019). Further, anecdotes regarding quicker rapport building with patients when working with a facility dog (Barnett, 2019) imply that the facility dog might allow for better connections and communications with pediatric patients. Many of the participants in the present study were child-life specialists whose job is to help hospitalized children cope with healthcare-related stress, pain, and fear in developmentally appropriate ways. It seems likely that faster rapport and better connections with patients, as catalyzed by the facility dog, would augment success in the role of a child-life specialist. This is consistent with findings that, after visiting with a therapy dog, pediatric patients described healthcare professionals as friendly and as making their hospital stays easier (Lindström Nilsson et al., 2020). On the other hand, there was no significant relationship between working with a facility dog and burnout in terms of perceived emotional exhaustion. Pediatric healthcare professionals working with facility dogs may still be seeing the same number of patients on a daily basis, if not more (Barnett, 2019). Therefore, it may be possible that these individuals experience just as much emotional exhaustion as their counterparts without facility dogs.

With respect to job-perceptions, working with a facility dog was significantly associated with more positive descriptions of one's job, greater job-related enthusiasm, and less job-related depression. Working with a facility dog was also associated with less intention to leave one's job and greater plans to stay in one's current position. This replicates similar findings from dogs in the workplace that suggest higher work-related morale (Foreman et al., 2017), better mood at work (Hall et al., 2017), and greater job satisfaction (R. T. Barker et al., 2012). Further, these findings may also be related to observations of less absenteeism and fewer sick days in workplaces where dogs are permitted (Wilkin et al., 2016). One explanation for these effects could be connected to lower stress levels at work when a dog is present (R. T. Barker et al., 2012). Specific to facility dogs, anecdotes from pediatric healthcare professionals have described feeling less anxious and being in a better mood when the facility dog is at work as well as feeling comforted by the facility dog's presence (Barnett, 2019). Finally, as pediatric healthcare professionals working with a facility dog were found to feel more accomplished at work, there could be connections between greater enthusiasm about going to work and lower chances of wanting to leave a job in which they are feeling accomplished.

There were no significant associations between working with a facility dog and reported

Accepted Article

feelings about coworkers or perceived support at work. This suggests that healthcare professionals with a facility dog do not feel any more positive about their coworkers than those who don't work with a facility dog. This finding is partially misaligned with the social support theory of human-animal interaction (Beetz, 2017; Wells, 2019), as perceived workplace support could have been an indicator for facility dogs acting as social catalysts between coworkers. This is also a stark contrast from several studies on dogs in the workplace that describe improved communication (R. T. Barker et al., 2012), greater social cohesion, and more positive interactions among coworkers in workplaces where dogs are present (Foreman et al., 2017). Within the healthcare setting, anecdotal reports on facility dogs have also described positive culture changes within the unit (Barnett, 2019) when dogs have been present. Although the present findings do not suggest that the social support theory is in action through improved coworker relations or hospital atmosphere, they do not rule out the possibility that the social support theory might be occurring through facility dogs providing direct companionship (Beetz, 2017; Wells, 2019).

Finally, working with a facility dog was significantly associated with better self-reported mental health in terms of more positive affect regarding recent experiences, less depression, less negative affect, and better perceptions of mental health. This suggests that pediatric healthcare professionals working with a facility dog may have better mental health than pediatric healthcare professionals without a facility dog in the form of feeling more positively overall. These findings are aligned with prior research suggesting that short interactions with a therapy dog can result in improved mood (Grajfoner et al., 2017), and fewer depressive symptoms (Souter & Miller, 2007). However, there was no significant relationship between working with a facility dog and self-reported anxiety. Based on extensive literature describing stress-reductions observed from human-dog interactions (S. B. Barker et al., 2010; Ein et al., 2018; Polheber & Matchock, 2014) this was an unexpected null finding. This could be in part due to the nature of the measure, as the PROMIS Anxiety scale is more commonly used to measure clinical levels of anxiety in response to targeted interventions. Because participants in the present sample did not score in the range of clinical anxiety, it is possible that the selected measure was not sensitive to specific sub-clinical changes within the sample. Further, this null finding could be related to some of the challenges described by pediatric healthcare professionals working with a facility dog. For example, one recurring theme in quotes about working with a facility dog is that the dog is always in high

Accepted Article

demand around the hospital. In response, the healthcare professional working with the facility dog must learn to manage frequently being stopped in the hallways and receiving more requests for the dog's presence than can be accommodated (Barnett, 2019). Anecdotes such as these suggest that working with a facility dog may add to the already high responsibilities of pediatric healthcare professionals, which might counteract any anxiety-reducing effects of human-animal interaction with the anxiety of higher work demands (e.g., Al Sabei et al., 2020; Buckley et al., 2020). It may be informative, in future research, to investigate sources of anxiety and make more specific distinctions between workplace anxiety and general anxiety.

Future Directions

Based on preliminary evidence for higher patient ratings of nursing communication after therapy dog interaction (Harper et al., 2015), it is speculated that the presence of a facility dog may also allow for greater rapport and improved interpersonal relationships with patients and their families. Evidence for this might suggest that the mechanism of social support in working with a facility dog takes place not through changes to pre-existing relationships with coworkers, but through new connections with patients and families. Thus, one future step in exploring the effects of facility dogs on healthcare professionals is to assess healthcare professionals', patients', and families' perceptions of provider-patient relationships. An assessment of provider-patient relationships may also reveal more about the difference in healthcare professionals' perceived accomplishment at work. As positive correlations have been found between pediatric nurses' perceptions of personal accomplishment and families' satisfaction with patient care (Buckley et al., 2020), a potential for better connections with patients and families when working with a facility dog may play a role in greater feelings of accomplishment.

Limitations

The present study provides early empirical evidence for the effects of facility dogs on pediatric healthcare professionals, but it is not without limitations that should be considered in future research. Data-specific limitations include low internal consistency (Cronbach's $\alpha = 0.33$) for SPANE Balanced Affect and the unsatisfied assumption of normality for JDI and JIG scales. The low internal consistency suggests that the SPANE may not have accurately measured affect balance in the sample, which could have created inaccurate regression estimates for this measure. The unsatisfied assumption of normality in small samples can bias estimates of standard error

Accepted Article

and heighten the risk of inflated significance in the results. However, in adequately-sized samples such as this, the risk of biased estimates in outcome transformations may be greater than risks associated with violations of the normality assumption (Schmidt & Finan, 2018). Thus, it is unlikely that the unsatisfied assumption of normality for JDI and JIG scales influenced their interpretations, but the possibility cannot be entirely ruled out. Limitations in the study design include the lack of randomized group assignment, which prevents results from being interpreted with any determination of causality. Additionally, there is likely to be participant self-selection bias within the sample. By recruiting from only one facility dog provider, training methods and organizational requirements are held constant within the sample, but generalizability is subsequently restricted. Generalizability is also limited by the inclusion of personnel in *pediatric* hospitals only, as there may be differing effects in adult hospitals or other healthcare settings. As most participants with facility dogs and control participants in the sample worked in separate hospitals (or at least in separate units), it was not possible to determine whether positive effects are related to the facility dogs themselves, or to broader hospital differences. For example, it is possible that some common characteristics of hospitals that house facility dog programs may be contributing to lower work-related burnout. Thus, it may be beneficial in future research to assess additional hospital-related measures such as absenteeism and staff performance as well as including a greater number of hospitals to investigate whether the present findings may be replicated or supported. Finally, changes over time cannot be determined from a cross-sectional design. Future longitudinal research is recommended to assess within-participant changes before and after being partnered with a facility dog.

Conclusion

In conclusion, findings from this project suggest the potential for facility dogs in pediatric hospitals to benefit healthcare professionals. This study found that, compared to the control group, those working with a facility dog reported less work-related burnout, more positive job perceptions, and better mental health. Specifically, benefits were observed in the form of greater perceived accomplishment, more positive feelings about work, less depression, and more positive affect overall. This provides information on the possible benefits of facility dog programs, such as staff-specific facility dog interventions for job-related well-being. The results of this project also indicated that working with a facility dog had no significant effects on emotional exhaustion, feelings about coworkers, perceived workplace support, and anxiety.

Accepted Article

Areas without significant group differences provide information on the boundaries of facility dog influences.

Relevance to Clinical Practice

Results indicated that working with a facility dog had significant positive effects on burnout, job-related perceptions, and mental health among pediatric healthcare professionals. This suggests potential value in facility dog interventions to improve staff job-related and personal wellbeing. Additionally, non-significant findings suggest the specificity of facility dog influences. These findings may inform hospitals with existing facility dog programs on how they can maximize the applicability and reach of their programs. Results may also inform hospitals considering the possibility of adding a facility dog program on what to expect, such that the anticipated role of a facility dog remain proportionate to the empirical evidence. Thus, observed effects from this study have the potential to benefit the healthcare industry through enhancing healthcare professionals' well-being while also setting realistic expectations for both existing and future facility dog programs.

References

- Al Sabei, S. D., Labrague, L. J., Miner Ross, A., Karkada, S., Albashayreh, A., Al Masroori, F., & Al Hashmi, N. (2020). Nursing Work Environment, Turnover Intention, Job Burnout, and Quality of Care: The Moderating Role of Job Satisfaction. *Journal of Nursing Scholarship*, 52(1), 95–104. <https://doi.org/10.1111/jnu.12528>
- Allen, K. M., Blascovich, J., Tomaka, J., & Kelsey, R. M. (1991). Presence of human friends and pet dogs as moderators of autonomic responses to stress in women. *Journal of Personality and Social Psychology*, 61(4), 582–589. <https://doi.org/10.1037/0022-3514.61.4.582>
- Barker, R. T., Knisely, J. S., Barker, S. B., Cobb, R. K., & Schubert, C. M. (2012). Preliminary investigation of employee's dog presence on stress and organizational perceptions. *International Journal of Workplace Health Management*, 5(1), 15–30. <https://doi.org/10.1108/17538351211215366>
- Barker, S. B., Knisely, J. S., McCain, N. L., & Best, A. M. (2005). Measuring stress and immune response in healthcare professionals following interaction with a therapy dog: A pilot study. *Psychological Reports*, 96(3), 713–729. <https://doi.org/10.2466/pr0.96.3.713-729>
- Barker, S. B., Knisely, J. S., McCain, N. L., Schubert, C. M., & Pandurangi, A. K. (2010). Exploratory study of stress-buffering response patterns from interaction with a therapy dog. *Anthrozoös*, 23(1), 79–91. <https://doi.org/10.2752/175303710X12627079939341>
- Barnett, B. N. (2019). *An exploration of child life programming and facility-canine assisted therapy in pediatric hospital settings* [Master's thesis, Missouri State University]. MSU Graduate Theses. 3398. <https://bearworks.missouristate.edu/theses/3398>
- Beetz, A. M. (2017). Theories and possible processes of action in animal assisted interventions. *Applied Developmental Science*, 21(2), 139–149. <https://doi.org/10.1080/10888691.2016.1262263>
- Boateng, G. O., Schuster, R. C., & Odei Boateng, M. (2019). Uncovering a health and wellbeing gap among professional nurses: Situated experiences of direct care nurses in two Canadian cities. *Social Science & Medicine*, 242, 112568. <https://doi.org/10.1016/j.socscimed.2019.112568>
- Bothma, C. F. C., & Roodt, G. (2013). The validation of the turnover intention scale. *SA Journal of Human Resource Management*, 11(1). <http://dx.doi.org/10.4102/sajhrm.v11i1.507>

- Buckley, L., Berta, W., Cleverley, K., Medeiros, C., & Widger, K. (2020). What is known about paediatric nurse burnout: A scoping review. *Human Resources for Health, 18*(1), 9. <https://doi.org/10.1186/s12960-020-0451-8>
- Carr, D. C., Taylor, M. G., Gee, N. R., & Sachs-Ericsson, N. (2020). Psychological health benefits of companion animals following a social loss. *The Gerontologist, 60*(3), 428–438. <https://doi.org/10.1093/geront/gnz109>
- Cella, D., Riley, W., Stone, A., Rothrock, N., Reeve, B., Yount, S., Amtmann, D., Bode, R., Buysse, D., Choi, S., Cook, K., DeVellis, R., DeWalt, D., Fries, J. F., Gershon, R., Hahn, E. A., Lai, J.-S., Pilkonis, P., Revicki, D., ... Hays, R. (2010). The Patient-Reported Outcomes Measurement Information System (PROMIS) developed and tested its first wave of adult self-reported health outcome item banks: 2005–2008. *Journal of Clinical Epidemiology, 63*(11), 1179–1194. <https://doi.org/10.1016/j.jclinepi.2010.04.011>
- Cohen, J. (1992). A power primer. *Psychological Bulletin, 112*(1), 155–159. <https://doi.org/10.1037/0033-2909.112.1.155>
- Crossman, M. K., Kazdin, A. E., Matijczak, A., Kitt, E. R., & Santos, L. R. (2020). The influence of interactions with dogs on affect, anxiety, and arousal in children. *Journal of Clinical Child & Adolescent Psychology, 49*(4), 535–548. <https://doi.org/10.1080/15374416.2018.1520119>
- Diener, E., Wirtz, D., Tov, W., Kim-Prieto, C., Choi, D., Oishi, S., & Biswas-Diener, R. (2010). New well-being measures: Short scales to assess flourishing and positive and negative feelings. *Social Indicators Research, 97*(2), 143–156. <https://doi.org/10.1007/s11205-009-9493-y>
- Ein, N., Li, L., & Vickers, K. (2018). The effect of pet therapy on the physiological and subjective stress response: A meta-analysis. *Stress and Health, 34*(4), 477–489. <https://doi.org/10.1002/smi.2812>
- Foreman, A., Glenn, M., Meade, B., & Wirth, O. (2017). Dogs in the workplace: A review of the benefits and potential challenges. *International Journal of Environmental Research and Public Health, 14*(5), 498. <https://doi.org/10.3390/ijerph14050498>
- Grajfoner, D., Harte, E., Potter, L., & McGuigan, N. (2017). The effect of dog-assisted intervention on student well-being, mood, and anxiety. *International Journal of*

Environmental Research and Public Health, 14(5), 483.
<https://doi.org/10.3390/ijerph14050483>

- Hall, S., Wright, H., McCune, S., Zulch, H., & Mills, D. (2017). Perceptions of dogs in the workplace: The pros and the cons. *Anthrozoös*, 30(2), 291–305. <https://doi.org/10.1080/08927936.2017.1311053>
- Harper, C. M., Dong, Y., Thornhill, T. S., Wright, J., Ready, J., Brick, G. W., & Dyer, G. (2015). Can therapy dogs improve pain and satisfaction after total joint arthroplasty? A randomized controlled trial. *Clinical Orthopaedics and Related Research®*, 473(1), 372–379. <https://doi.org/10.1007/s11999-014-3931-0>
- Hays, R. D., Schalet, B. D., Spritzer, K. L., & Cella, D. (2017). Two-item PROMIS® global physical and mental health scales. *Journal of Patient-Reported Outcomes*, 1. <https://doi.org/10.1186/s41687-017-0003-8>
- Ironson, G., Smith, P., Brannick, M., Gibson, W., & Paul, K. (1989). Construction of a job in general scale: A comparison of global, composite, and specific measures. *Journal of Applied Psychology - J APPL PSYCHOL*, 74, 193–200. <https://doi.org/10.1037/0021-9010.74.2.193>
- Kim, H., & Kao, D. (2014). A meta-analysis of turnover intention predictors among U.S. child welfare workers. *Children and Youth Services Review*, 47, 214–223. <https://doi.org/10.1016/j.childyouth.2014.09.015>
- Kline, J. A., VanRyzin, K., Davis, J. C., Parra, J. A., Todd, M. L., Shaw, L. L., Haggard, B. R., Fisher, M. A., Pettit, K. L., & Beck, A. M. (2020). Randomized trial of therapy dogs versus deliberative coloring (art therapy) to reduce stress in emergency medicine providers. *Academic Emergency Medicine*, 27(4), 266–275. <https://doi.org/10.1111/acem.13939>
- Laguna, M., Mielniczuk, E., Razmus, W., Moriano, J. A., & Gorgievski, M. J. (2017). Cross-culture and gender invariance of the Warr (1990) job-related well-being measure. *Journal of Occupational and Organizational Psychology*, 90(1), 117–125. <https://doi.org/10.1111/joop.12166>
- Lee, E., Daugherty, J., Eskierka, K., & Hamelin, K. (2019). Compassion Fatigue and Burnout, One Institution's Interventions. *Journal of Perianesthesia Nursing: Official Journal of*

the American Society of PeriAnesthesia Nurses, 34(4), 767–773.
<https://doi.org/10.1016/j.jopan.2018.11.003>

- Lindström Nilsson, M., Funkquist, E., Edner, A., & Engvall, G. (2020). Children report positive experiences of animal-assisted therapy in paediatric hospital care. *Acta Paediatrica*, 109(5), 1049–1056. <https://doi.org/10.1111/apa.15047>
- Maslach, C., Jackson, S., & Leiter, M. (1997). The Maslach Burnout Inventory Manual. In *Evaluating Stress: A Book of Resources* (Vol. 3, pp. 191–218).
- Mowday, R. T., Koberg, C. S., & McArthur, A. W. (1984). The psychology of the withdrawal process: A cross-validated test of Mobley's Intermediate Linkages Model of turnover in two samples. *The Academy of Management Journal*, 27(1), 79–94. JSTOR. <https://doi.org/10.2307/255958>
- Ortega-Campos, E., Cañadas-De la Fuente, G. A., Albendín-García, L., Gómez-Urquiza, J. L., Monsalve-Reyes, C., & de la Fuente-Solana, E. I. (2019). A Multicentre Study of Psychological Variables and the Prevalence of Burnout among Primary Health Care Nurses. *International Journal of Environmental Research and Public Health*, 16(18), 3242. <https://doi.org/10.3390/ijerph16183242>
- Polheber, J. P., & Matchock, R. L. (2014). The presence of a dog attenuates cortisol and heart rate in the Trier Social Stress Test compared to human friends. *Journal of Behavioral Medicine*, 37(5), 860–867. <https://doi.org/10.1007/s10865-013-9546-1>
- Powell, L., Edwards, K. M., McGreevy, P., Bauman, A., Podberscek, A., Neilly, B., Sherrington, C., & Stamatakis, E. (2019). Companion dog acquisition and mental well-being: A community-based three-arm controlled study. *BMC Public Health*, 19(1), 1428. <https://doi.org/10.1186/s12889-019-7770-5>
- Schalet, B. D., Pilkonis, P. A., Yu, L., Dodds, N., Johnston, K. L., Yount, S., Riley, W., & Cella, D. (2016). Clinical validity of PROMIS Depression, Anxiety, and Anger across diverse clinical samples. *Journal of Clinical Epidemiology*, 73, 119–127. <https://doi.org/10.1016/j.jclinepi.2015.08.036>
- Schmidt, A. F., & Finan, C. (2018). Linear regression and the normality assumption. *Journal of Clinical Epidemiology*, 98, 146–151. <https://doi.org/10.1016/j.jclinepi.2017.12.006>
- Shilp, J. (2019). *Dogs in Health Care: Pioneering Animal-Human Partnerships*. McFarland.

- Smith, P. C., Kendall, L. M., & Hulin, C. L. (1969). *The measurement of satisfaction in work and retirement: A strategy for the study of attitudes* (pp. xiii, 186). Rand McNally.
- Souter, M. A., & Miller, M. D. (2007). Do animal-assisted activities effectively treat depression? A meta-analysis. *Anthrozoös*, *20*(2), 167–180. <https://doi.org/10.2752/175303707X207954>
- Spruin, E., Dempster, T., & Mozova, K. (2020). Facility dogs as a tool for building rapport and credibility with child witnesses. *International Journal of Law, Crime and Justice*, 100407. <https://doi.org/10.1016/j.ijlcj.2020.100407>
- Tavakol, M., & Dennick, R. (2011). Making sense of Cronbach's alpha. *International Journal of Medical Education*, *2*, 53–55. <https://doi.org/10.5116/ijme.4dfb.8dfd>
- Tawfik, D. S., Profit, J., Morgenthaler, T. I., Satele, D. V., Sinsky, C. A., Dyrbye, L. N., Tutty, M. A., West, C. P., & Shanafelt, T. D. (2018). Physician Burnout, Well-being, and Work Unit Safety Grades in Relationship to Reported Medical Errors. *Mayo Clinic Proceedings*, *93*(11), 1571–1581. <https://doi.org/10.1016/j.mayocp.2018.05.014>
- Velando-Soriano, A., Ortega-Campos, E., Gómez-Urquiza, J. L., Ramírez-Baena, L., Fuente, E. I. D. L., & Fuente, G. A. C.-D. L. (2020). Impact of social support in preventing burnout syndrome in nurses: A systematic review. *Japan Journal of Nursing Science*, *17*(1), e12269. <https://doi.org/10.1111/jjns.12269>
- von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gøtzsche, P. C., & Vandenbroucke, J. P. (2008). STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Journal of Clinical Epidemiology* *61*(4), 344–3499. PMID: 18313558
- Walsh, D., Yamamoto, M., Willits, N. H., & Hart, L. A. (2018). Job-related stress in forensic interviewers of children with use of therapy dogs compared with facility dogs or no dogs. *Frontiers in Veterinary Science*, *5*, 46. <https://doi.org/10.3389/fvets.2018.00046>
- Warr, P. (1990). The measurement of well-being and other aspects of mental health. *Journal of Occupational Psychology*, *63*(3), 193–210. <https://doi.org/10.1111/j.2044-8325.1990.tb00521.x>
- Wei, H., Kifner, H., Dawes, M. E., Wei, T. L., & Boyd, J. M. (2020). Self-care Strategies to Combat Burnout Among Pediatric Critical Care Nurses and Physicians. *Critical Care Nurse*, *40*(2), 44–53. <https://doi.org/10.4037/ccn2020621>

- Wells, D. L. (2019). The state of research on human–animal relations: Implications for human health. *Anthrozoös*, 32(2), 169–181. <https://doi.org/10.1080/08927936.2019.1569902>
- Wilkin, C. L., Fairlie, P., & Ezzedeen, S. R. (2016). Who let the dogs in? A look at pet-friendly workplaces. *International Journal of Workplace Health Management*, 9(1), 96–109. <https://doi.org/10.1108/IJWHM-04-2015-0021>
- Wood, L. J., Giles-Corti, B., Bulsara, M. K., & Bosch, D. A. (2007). More than a furry companion: The ripple effect of companion animals on neighborhood interactions and sense of community. *Society & Animals*, 15(1), 43–56. <https://doi.org/10.1163/156853007X169333>
- Yoon, J., & Thye, S. (2000). Supervisor support in the work place: Legitimacy and positive affectivity. *The Journal of Social Psychology*, 140, 295–316. <https://doi.org/10.1080/00224540009600472>
- Yordy, M., Brandon, A., & Harmon, L. (2020). Animal-assisted therapy: Promoting a healthy work environment. *The Journal of Excellence in Nursing Leadership*, 51(4), 24–30. <https://doi.org/10.1097/01.NUMA.0000657248.63754.fc>

Table 1

Sample characteristics

	Control (<i>n</i> = 65)		Facility Dog (<i>n</i> = 65)		Group Difference	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>t</i>	<i>p</i>
Age	37.53	8.97	37.48	10.44	–	–
Time in job position (years)	4.25	4.45	5.08	4.76	-1.017	.311
	<i>n</i>	%	<i>n</i>	%	χ^2	<i>p</i>
Gender identity					–	–
Male	5	7.69	5	7.69		
Female	60	92.31	60	92.31		
Relationship status					8.013	.155
Married	31	47.69	41	63.07		
Living with partner	7	10.77	4	6.15		
Single, never married	17	26.15	18	27.69		
Separated	2	3.08	0	0.00		
Divorced	7	10.77	2	3.08		
Widowed	1	1.54	0	0.00		
Race/Ethnicity					7.203	.206
White	59	90.77	64	98.46		
Hispanic or Latino	2	3.08	0	0.00		
Black or African American	2	3.08	0	0.00		
Asian Indian	2	3.08	0	0.00		
American Indian or Alaska Native	0	0.00	1	1.54		

Children					–	–
Yes	29	44.62	29	44.62		
No	36	55.38	36	55.38		
Pet dog					0.034	.854
Yes	42	64.62	43	66.15		
No	23	35.38	22	33.85		

Note. Group difference test statistics are omitted for characteristics matched across groups (age, gender identity) and for unmatched characteristics found to be identical across groups (children). In all cases where a participant selected multiple options for race/ethnicity, the selections were for one minority race/ethnicity and white. These participants are recorded once in the table, each under their selected minority race/ethnicity. No participants reported gender identity in the category of other or prefer not to answer. No participants reported race/ethnicity in the categories of Asian/Native Hawaiian or Pacific Islander/prefer not to answer.

Table 2

Measure characteristics, Cronbach's alpha in the sample, and group differences

	Items	Scoring	Cronbach's α	Control	Facility Dog	Cohen's d
				$M (SD)$	$M (SD)$	
Work-related burnout						
MBI: Emotional Exhaustion	9	Sum: 0-54	0.93	42.63 (10.79)	43.52 (10.84)	0.08
MBI: Personal Accomplishment	8	Sum: 0-48	0.79	46.02 (6.42)	50.75 (3.65)	0.91
Job perceptions						
Job-Related Depression and Enthusiasm	6	Sum: 6-30	0.85	25.74 (3.90)	27.40 (3.00)	0.48
JDI: People in Your Present Job	18	Sum: 0-54	0.88	43.25 (8.76)	45.00 (7.06)	0.22
Job in General	18	Sum: 0-54	0.81	46.45 (9.78)	50.86 (4.86)	0.57

Workplace Social Support	9	Average	0.83	3.65 (0.70)	3.87 (0.69)	0.31
Anticipated Turnover	12	Average	0.87	4.74 (1.37)	5.32 (1.07)	0.47
Turnover Intention	6	Average	0.87	2.65 (0.79)	2.28 (0.67)	-0.51
Mental health						
PROMIS: Depression	8	Manual T-score	0.93	48.75 (7.78)	45.80 (6.84)	-0.40
PROMIS: Anxiety	8	Manual T-score	0.89	54.38 (7.11)	53.02 (6.22)	-0.20
PROMIS: Global Mental Health	2	Manual T-score	0.75	38.00 (6.16)	34.94 (6.96)	-0.47
SPANE: Positive Affect	6	Sum: 6-30	0.84	23.26 (3.94)	25.46 (3.15)	0.62
SPANE: Negative Affect	6	Sum: 6-30	0.91	14.66 (3.95)	13.43 (3.27)	-0.34
SPANE: Balanced Affect	12	Sum: -24 to 24	0.33	8.60 (7.21)	12.03 (5.53)	0.53

Note. MBI = Maslach Burnout Inventory. JDI = Job Descriptive Index. PROMIS = Patient Reported Outcomes Measurement Information System. SPANE = Scale of Positive and Negative Experience. Cronbach's α shows good reliability for a measure if it is greater than 0.8, and adequate reliability for a measure if it is greater than 0.7 (Tavakol & Dennick, 2011).

Table 3

Hierarchical regression for cross-sectional group comparison of participants with facility dogs versus matched controls

	Model 1:		Model 2: Demographics + Facility Dog					
	Demographics		<i>F</i>	<i>R</i> ²	Specific Effects of Facility Dog			
	<i>F</i>	<i>R</i> ²			β	95% CI	ΔF	ΔR^2
Work-related burnout								
MBI: Emotional Exhaustion	4.70***	0.21	4.15***	0.22	0.05	-2.35 to 4.68	0.43	0.00
MBI: Personal Accomplishment	0.30	0.02	3.34**	0.18	0.42***	2.83 to 6.65	24.26***	0.16
Job perceptions								
Job-Related Depression and Enthusiasm	2.58*	0.13	3.43***	0.19	0.24**	0.53 to 2.91	8.25**	0.06
JDI: People in Your Present Job	1.46	0.08	1.50	0.09	0.12	-0.96 to 4.64	1.69	0.01
Job in General	2.29*	0.12	3.23**	0.18	0.25**	1.33 to 6.68	8.76**	0.06
Workplace Social Support	2.03	0.10	2.21*	0.13	0.16	-0.02 to 0.46	3.25	0.02
Anticipated Turnover	1.35	0.07	2.23*	0.13	0.24**	0.18 to 1.05	7.87**	0.06
Turnover Intention	1.64	0.09	2.77**	0.16	-0.27**	-0.66 to -0.15	9.89**	0.07
Mental health								
PROMIS: Depression	2.08*	0.11	2.52**	0.14	-0.20*	-5.44 to -0.37	5.13*	0.04
PROMIS: Anxiety	2.66**	0.13	2.56**	0.15	-0.12	-3.81 to 0.75	1.77	0.01
PROMIS: Global Mental Health	2.76**	0.14	3.24**	0.18	-0.21*	-5.00 to -0.50	5.84*	0.04
SPANE: Positive Affect	3.67***	0.17	5.21***	0.26	0.29***	1.00 to 3.36	13.37***	0.08
SPANE: Negative Affect	4.02***	0.19	4.22***	0.22	-0.18*	-2.51 to -0.12	4.75*	0.03

SPANE: Balanced Affect	4.68***	0.21	5.82***	0.28	0.27***	1.42 to 5.57	11.13***	0.07
------------------------	---------	------	---------	------	---------	--------------	----------	------

Note. MBI = Maslach Burnout Inventory. JDI = Job Descriptive Index. PROMIS = Patient Reported Outcomes Measurement Information System. SPANE = Scale of Positive and Negative Experience. CI = Confidence Interval. For Block 1 of each model $df = (7, 122)$ and for Block 2 of each model $df = (1, 121)$. $N = 130$. ΔF is the primary metric for determining whether there was a difference in the outcome measure uniquely related to facility dog presence or absence, above and beyond any difference that was related to participant demographics. $*p \leq .05$, $**p \leq .01$, $***p \leq .001$